GENERAL INDEX

New scientific names of plants and the final members of new combinations are printed in **bold face** type; synonyms and page numbers having reference to figures or plates, in *italic;* and previously published scientific names and all other matter, in ordinary type.

A

Acid, oxalic, produced by Sclerotinia cinerea, 319
Acidity: The effect of certain conditions upon the, of tomato fruits, 229; of plum fruits, 317; relation of growth of Sclerotinia cinerea to, 318
Agaricaceæ, 197
Agaricus betulinus, 146; *quercinus*, 145
Air, The identification of the most characteristic salivary organism, and its relation to the pollution of, 47
Air-sampling apparatus, bacteriological, *80;* construction and use of, 66
albido-brunnea (Thelephora), 214, 216, *228*
albo-violascens (Cyphella), 364
Aleurodiscus, 198
Algæ: A contribution to our knowledge of the relation of certain species of grass-green, to elementary nitrogen, 157; Some pure culture methods in the, 23; Indications regarding the source of combined nitrogen for Ulva Lactuca, 229
angustata (Thelephora), 205
anomala (Solenia), 373, var. *arbicularis*, 373
Anthericum **Chandleri,** 405; stenocarpum, 406
anthocephala (Thelephora), 202, 203, *226*
arachnoidea (Cyphella), 363
Arrhenia, 344
artocreas (Michenera), 197
Asterostroma, 198
Augustinii (Heterothecium), 381

B

Bananæ (Cyphella), 379
biennis (Thelephora), 215
Boletus abietinus, 91; *adustus*, 102; *applanatus*, 137; *arcularius*, 107; *betulinus*, 104; *brumalis*, 107; *caesius*, 96; *cinnabarinus*, 116; *cinnamomeus*, 123; *conchatus*, 132; *conchifer*, 93; *confragosus*, 144; *cuticularis*, 118; *distortus*, 105; *dryadeus*, 119; *elegans*, 110; *fomentarius*, 136; *fraxineus*, 130; *frondosus*, 112; *fulvus*, 133; *fumosus*, 103; *gilvus*, 117; *giganteus*, 113; *graveolens*, 131; *hirsutus*, 93; *hispidus*, 119; *igniarius*, 135; *lucidus*, 123; *nummularius*, 110; *perennis*, 122; *Pini*, 142; *pinicola*, 130; *pubescens*, 94; *radiatus*, 118; *resinosus*, 116; *sanguineus*, 115; *spumeus*, 99; *squamosus*, 109; *suaveolens*, 140; *sulphureus*, 114; *tulipifera*, 152; *umbellatus*, 112; *unicolor*, 143; *versicolor*, 91
Bordeaux mixture, effect of, on transpiration, 12, 351
borealis (Craterellus), 357, *382*
Botrydiopsis sp., isolation of, in pure culture, 38
Botrydium granulatum, isolation of, in pure culture, 38
brevipes (*Cantharellus*), 329
Burt, E. A. The Thelephoraceæ of North America, I, 185; II, 327; III, 357

C

Cacalia prenanthoides, 271; *runcinata*, 272; *Toluccana*, 271
caespitulans (Thelephora), 204, *346*
calceum (Corticium), 187
calyculus (Craterellus), 338
campanula (*Peziza*), 360
canadensis (Cantharellus), 345
canadensis (Craterellus), 345
candida (*Cyphella*), 377
Cantharellus, 197; *brevipes*, 329; canadensis, 345; floccosus, 345; *lævis*, 362
Cantharellus (Craterellus), 330, 332, *346*
capula (Cyphella), 366, *382*
caricina (Cyphella), 366, *382*
caryophyllea (Thelephora), 207, 209, *226*
Castor bean leaves, use of, in potometer experiments, 7, 11

Cellulose, methods of preparation of, 305
Chlamydomonas pisiformis forma minor, isolation of, in pure culture, 32; relation of, to elementary nitrogen, 178, 179, 181
Chlorascens (Heterobasidium), 197
Chlorella sp., and C. vulgaris, isolation of, in pure culture, 33; relation of, to elementary nitrogen, 178, 179, 181
Chlorococcum humicola, isolation of, in pure culture, 35; relation of, to elementary nitrogen, 178, 179, 181
cinereo-fusca (Cyphella), 377
cinereo-fusca (*Peziza*), 377
Cladoderris, 198
Clavaria pistillaris, 342
clavatus (Craterellus), 329, 345, *346*
confluens (*Craterellus*), 332
conglobata (Cyphella), 375, *382*
Coniophora, 198, 199
convoluta (Cyphella), 380
Cooley, J. S.: A study of the physiological relations of Sclerotinia cinerea (Bon.) Schröter, 291; Duggar, B. M., and, The effect of surface films and dusts on the rate of transpiration, 1, The effects of surface films on the rate of transpiration: experiments with potted potatoes, 351
Cora, 199
corbiformis (Thelephora), 211
Coriolus Lloydii, 95; *nigromarginatus*, 93
cornucopioides (Craterellus), 327, 333, *350*
cornucopioides (Thelephora), 212, *346*
corrugis (*Craterellus*), 340
Corticium, 190, 191, 192, 193, 197, 198, 199; calceum, 187; lactescens, latex in, 194; lacteum, 187; Sambuci, 191; subgiganteum, 197; vagum, 197
Craterellus, 190, 196, 197, 327; **borealis,** 357, *382;* **calyculus,** 338; *canadensis,* 345; Cantharellus, 330, 332, *346;* clavatus, 329, 345, *346; confluens,* 332; cornucopioides, 327, 333, *350; corrugis,* 340; crispus, 338; **delitescens,** 339, *350;* **dilatus,** 343, *350;* dubius, 335; **Humphreyi,** 344, *350; lateritius,* 330; lutescens, 336, *350; Pogonati,* 362, *382;* marasmioides, 345; **ochrosporus,** 327, 334, *350;* ocreatus, 334; odoratus, 327, 331, *346, 348;* **palmatus,** 342, *350;* pistillaris, 327, 341, *348, 350; Pogonati,* 362, *382;* pulverulentus, 345; roseus, 332; sinuosus, 337; spathularius, 345; taxophilus, 339, *350;* unicolor, 327, 340, 341, 342, *348*
crispus (Craterellus), 338
cupulæformis (Cyphella), 369, *382*
Cupressi (Cyphella), 380
cuticularis (Thelephora), 216, *228*
Cyclomyces, 82, 147; Greenei, 147
Cymatella, 345
Cyphella, 197, 358; albo-violascens, 364; arachnoidea, 363; Bananæ, 379; *candida,* 377; capula, 366, *382;* caricina, 366, *382;* cinereo-fusca, 377; **conglobata,** 375, *382;* convoluta, 380; Cupressi, 380; cupulæformis, 369, *382;* digitalis, 358; fasciculata, 373, 376, *382;* filicicola, 379; fraxinicola, 368; *fulva,* 373; fumosa, 376, *382; furcata,* 373; galeata, 362, *382;* globosa, 367; griseopallida, 369; læta, 361; **Langloisii,** 368, *382;* **mellea,** 372, *382;* **minutissima,** 367, 368, *382;* musæcola, 380; muscigena, 362, 363, *382;* Palmarum, 377; Peckii, 377; perexigua, 378; pezizoides, 364, 365, 378; **porrigens,** 368, *382;* punctiformis, 367; Ravenelii, 371, 372, 373, *382; Ravenelii* Sacc., 373; *Saccardoi,* 373; *subcyanea,* 380; subgelatinosa, 370; sulphurea, 360; texensis, 371, 373, *382;* Tiliæ, 364, 365, *382;* trachychæta, 379; villosa, 365, *382*

D

Daedalea, 82, 143; ambigua, 144; confragosa, 144; *mollis,* 141; *pallido-fulva,* 146; quercina, 145; *sæpiaria,* 147; unicolor, 143
Davis, A. R., Duggar, B. M. and. Enzyme action in Fucus vesiculosus, 419
delitescens (Craterellus), 339, *350*
Dendrocladium, 199
dentosa (Thelephora), 224, *346*
digitalis (Cyphella), 358
dilatus (Craterellus), 343, *350*
dubius (Craterellus), 335
Duggar, B. M.: and Cooley, J. S., The effect of surface films and dusts on the rate of transpiration, 1, The effects of surface films on the rate of transpiration: experiments with potted potatoes, 351; and Davis, A. R., Enzyme action in Fucus vesiculosus, 419; and Merrill, M. C., The effect of certain conditions upon the acidity of tomato fruits, 229
Dusts, the effect of surface films and, on the rate of transpiration, 1

E

Enslinia pocula, 106
Enzymes: cellulase in Sclerotinia cinerea and Penicillium expansum, 305; pectinase in Sclerotinia cinerea and Penicillium expansum, 312
Erechthites runcinata, 271, 272
Erinus tomentosus, 409
Eu-Thelephoreæ, 197
Exobasidium, 193, 197

F

fasciculata (Cyphella), 373, 376, *382*
Favolus, 82, 148; canadensis, 148; *ohiensis*, 148; rhipidium, 148; *striatulus*, 148
filicicola (Cyphella), 379
Films and dusts, the effect of surface, on the rate of transpiration, 1
Films, the effect of surface, on the rate of transpiration: experiments with potted potatoes, 351
fimbriata (Thelephora), 222, *226*
Flaveria longifolia, 413, var. **subtomentosa**, 413
Flowering plants, Diagnoses of, chiefly from the southwestern United States and Mexico, 405
Fomes, 82, 126; applanatus, 130, 135, 136, 137, 138; carneus, 131; conchatus, 132; connatus, 129; Everhartii, 134, 135; fomentarius, 135, 136, 137; fraxineus, 129, 130; fraxinophilus, 129; fulvus, 133; graveolens, 131; igniarius, 134, 135; lobatus, 137; nigricans, 134, 135, 136; ohiensis, 128; pinicola, 130; *populinus*, 130; ribis, 133; rimosus, 133, cause of trunk disease of mesquite, 248; roseus, 131; *salicinus*, 133; scutellatus, 128; *supinus*, 133
Foster, G. L. Indications regarding the source of combined nitrogen for Ulva Lactuca, 229
fraxinicola (Cyphella), 368
Fucus vesiculosus, Enzyme action in, 419
fulva (*Cyphella*), 373
fumosa (Cyphella), 376, *382*
Fungous and host cells, A method for the differential staining of, 241.
furcata (*Cyphella*), 373

G

galeata (Cyphella), 362, *382*
Ganoderma sessile, 123; *subperforatum*, 123
Gates, R. R. A Texan species of Megapterium, 401; Some œnotheras from Cheshire and Lancashire, 383
globosa (Cyphella), 367
Gloeocystidium, 193
Gloeopeniophora, 193
Gloeoporus, 82, 149; conchoides, 149
Grandinia, 196
Greenman, J. M.: Descriptions of North American Senecioneæ, 263; and Thompson, C. H., Diagnoses of flowering plants, chiefly from the southwestern United States and Mexico, 405
Grifola poripes, 111; *ramosissima*, 112; *Sumstinei*, 113
griseo-pallida (Cyphella), 369
griseozonata (Thelephora), 221, *228*

H

Herpestes tomentosa, 409
Heterobasidium, 197; chlorascens, 197
Heterothecium Augustinii, 381
Hirneolina, 197, 198
hiscens (*Thelephora*), 207
Humphreyi (Craterellus), 344, *350*
Hydnaceæ, 196
Hymenochaete, 193, 198
Hymeno-lichens, 199
Hyphomycetes, 194
Hypochnaceæ, 190
Hypochnus, 190, 193, 196, 198, 199
Hypolyssus, 198

I

Infection and penetration of plums by Sclerotinia cinerea, 297
Inonotus texanus, 247
Introduction, i
intybacea (Thelephora), 217, 220, *228*
Irpex, 82, 151; cinnamomeus, 152; farinaceus, 152, 153; mollis, 152; tulipifera, 152

K

Kirchneriella sp., isolation of, in pure culture, 34; relation of, to elementary nitrogen, 178, 179, 181
Kummeria, 409

L

Lachnocladium, 196, 198, 199
laciniata (*Thelephora*), 219
laciniatum (*Stereum*), 219
lactescens (Corticium), 194
lacteum (Corticium), 187
læta (Cyphella), 361
lævis (*Cantharellus*), 362
Langloisii (Cyphella), 368, *382*
lateritius (*Craterellus*), 330
Lenzites, 82, 145; betulina, 146; *Crataegi*, 144; *flaccida*, 146; protractus, destruction of sap-wood of mesquite by, 248; saepiaria, 147; vialis, 146
Lilac, A trunk disease of the, 253
Lloydella, 192
lutescens (Craterellus), 336, *350*
lutosa (Thelephora), 216, *346*

M

magnispora (Thelephora), 211, *226*
Mappia, 408
marasmioides (Craterellus), 345

Megapterium, A Texan species of, 401; **argyrophyllum**, 401, *404*, var. **retusifolium**, 401, *404*; Fremontii, 402; **macrocarpum**, 402; missouriensis, 402, 403
mellea (Cyphella), 372, *382*
Merrill, M. C., Duggar, B. M., and. The effect of certain conditions upon the acidity of tomato fruits, 229
Merulius, 82, 150, 197; lacrymans, 151; rubellus, 150; tremellosus, 151; *tremellosus*, 151
Mesquite, Two trunk diseases of the, 243
Michenera, 197; artocreas, 197
Micheneri (*Stereum*), 214
Microcoleus sp., isolation of, in pure culture, 39
minutissima (Cyphella), 367, 368, *382*
Moore, G. T. Foreword, i
Mucronoporus Everhartii, 134
multipartita (Thelephora), 205, 208, *226*
musæcola (Cyphella), 380
muscigena (Cyphella), 362, *382*
Mycobonia, 198

N

Navicula sp., isolation of, in pure culture, 39
Nitrogen: A contribution to our knowledge of the relation of certain species of grass-green algæ to elementary, 157; Indications regarding the source of combined, for Ulva Lactuca, 229
Nolte, A. G. The identification of the most characteristic salivary organism, and its relation to the pollution of air, 47

O

obovata (Xylaria), 225
ochrosporus (Craterellus), 334, *350*
ocreatus (Craterellus) 334, 335
odoratus (Craterellus), 331, *346*, *348*
odorifera (*Thelephora*), 215
Oecopetalum, 408; **mexicanum**, 408, *416*
Oedogonium, isolation of zoospores free from bacteria in, 36
Oenothera Lamarckiana, 393, *400*; *macrocarpa*, 402; multiflora, 386, *396*; var. **elliptica**, 387, *396*, *398*; **rubrinervoides**, 389, *396*, *398*; **rubritincta**, 391, *398*, *400*; **tardiflora**, 391, *400*
Oenotheras, Some, from Cheshire and Lancashire, 383
Ohio, The Polyporaceæ of, 81
Oscillatoria sp., isolation of, in pure culture, 39
Overholts, L. O. The Polyporaceæ of Ohio, 81

P

palmata (Thelephora), 201, *226*
palmata americana (*Thelephora*), 201
Palmarum (Cyphella), 377
palmatus (Craterellus), 342, *350*
pannosa (Thelephora), 207
Peckii (Cyphella), 377
Penicillium expansum, physiological studies in, 308, 315, 322
Peniophora, 190, 191, 192, 193, 198, 199
Peniophorella, 193
perexigua (Cyphella), 378
Permeability of collodion membranes, 40
perplexa (Thelephora), 223, *346*
Peziza *campanula*, 360; *Tiliæ*, 364
pezizoides (Cyphella), 364, 365, 378
pistillaris (Clavaria), 342
pistillaris (Craterellus), 341, *348*, *350*
Pleurococcus vulgaris, isolation of, in pure culture, 34
Pogonati (*Craterellus*), 362, *382*
Polyporaceæ, 197
Polyporaceæ of Ohio, The, 81; index to species of, 153; key to genera of, 85
Polyporus, 82, 86; abietinus, 91, 92; *abortivus*, 105; adustus, 102, 103, 104; *albellus*, 97; *anax*, 113, 114; *applanatus*, 137; *arculariformis*, 107; arcularius, 107; *badius*, 126; Berkeleyi, 113; betulinus, 104; biformis, 95; borealis, 100; brumalis, 107; caesius, 96, 97; *carneus*, 131; *castanophilus*, 115; chioneus, 97; *cincinnatus*, 114; cinnabarinus, 116; cinnamomeus, 123; circinatus, 121, 122, 123; *conchatus*, 132; conchifer, 93; *conglobatus*, 131; *connatus* Fries, 129; *connatus* Schw., 122; *connatus* Weinm., 129; *cristatus*, 111; Curtisii, 125; cuticularis, 117, 118, 119; delectans, 99; *dichrous*, 150; distortus, 105; dryadeus, 119, 120; dryophilus, 119, 120; *dualis*, 121; elegans, 110; *endocrocinus*, 119; *fibula*, 95; *fissus*, 109; flavovirens, 111; focicola, 122, 123; *fomentarius*, 136; *fragilis*, 126; fragrans, 102, 103; *fraxineus*, 130; *fraxinophilus*, 129; frondosus, 112, 113, 114, 126; *fulvus*, 133; fumosus, 102, 103; galactinus, 97, 98, 141; giganteus, 113, 126; gilvus, 117, 118, 120; guttulatus, 100; hirsutulus, 91, 92; hirsutus, 93; hispidus, 119; *hypococcineus*, 115; *igniarius*, 135; *immitus*, 98; *intybaceus*, 126; *isidioides*, 117; lacteus, 97; *lentus*, 126; *leucomelas*, 126; *leucophæus*, 137; *lobatus*, 137; Lloydii, 95; lucidus, 123; *maculatus*, 100; *molliusculus*, 95, 96; *Morgani*, 110; nidulans, 117; *nigricans*, 136; obesus, 121, 123; obtusus, 100; *ovinus*, 126; pargamenus, 92; *parvulus*, 122; pennsylvanicus, 108; perennis, 122,

123; *perplexus*, 118; *phaeoxanthus*, 126; picipes, 109, 110; Pilotæ, 115; *pinicola*, 130; pocula, 106; *populinus*, 141; *puberula*, 103; pubescens, 94, 96, var. *Grayii*, 95; radiatus, 117, 118, 119; radicatus, 110; *reniformis*, 137; resinosus, 116; *rimosus*, 133; **robiniophila**, 104; Rostkowii, 109; *rufescens*, 106; sanguineus, 115, 116; Schweinitzii, 120; *scutellatus*, 128; semipileatus, 96; *serialis*, 139; Spraguei, 101; spumeus, 99; squamosus, 109; *subsericeus*, 123; *Sullivantii*, 94; sulphureus, 114; texanus, 247, cause of trunk disease of mesquite, 246, *250*, *252;* tomentosus, 121; umbellatus, 112, 113; varius, 110; *velutinus*, 93, 95; versicolor, 91, 93, cause of trunk disease of lilac, 253, *260*, *262; virgineus*, 94; volvatus, 105; zonalis, 101; *zonatus*, 91

Polystictus, 82; Lindheimeri, destruction of sap-wood of mesquite by, 248; *obesus*, 121; *proliferus*, 123

Poraqueiba, 409

Poria, 82

porrigens (Cyphella), 368, *332*

Potato plants, transpiration of, 351

Potometer, *22;* experiments with, 7

Prosopis glandulosa, trunk diseases of, 243

Protosiphon botryoides, isolation of, in pure culture, 35; relation of, to elementary nitrogen, 178, 179, 181

pulverulentus (Craterellus), 345

punctiformis (Cyphella), 367

Pure culture methods in the Algæ, Some, 23

pusiola (Thelephora), 208, var. terrestris, 209

puteana (Coniophora), 197

Pyropolyporus robiniæ, 133

R

radiata (Thelephora), 210

Randia aculeata, 411; **Gaumeri**, 410; **Purpusii**, 410; **truncata**, 411, *418;* xalapensis, 411

Ravenelii (Cyphella), 371, 372, *332*

Ravenelii (*Cyphella*), 373

Ravenelii (*Thelephora*), 207

regularis (Thelephora), 206, 207, *226*

Rhipidonema, 199

roseus (Craterellus), 332

Ruellia Parryi, 410

S

Saccardoi (*Cyphella*), 373

Salivary organism, The identification of the most characteristic, and its relation to the pollution of air, 47

Sambuci (Corticium), 191

sanguinolentum (Stereum), 194

Scenedesmus sp., isolation of, in pure culture, 34

Schizophyllum commune, destruction of sap-wood of mesquite by, 248

Schramm, J. R. A contribution to our knowledge of the relation of certain species of grass-green algæ to elementary nitrogen, 157; Some pure culture methods in the algæ, 23

von Schrenk, Hermann. A trunk disease of the lilac, 253; Two trunk diseases of the mesquite, 243

scissilis (Thelephora), 204, *226*

Sclerocarpus, **elongatus**, 412; Schiedeanus, 412; *Schiedeanus* var. *elongatus*, 412

Sclerotinia cinerea (Bon.) Schröter, A study of the physiological relations of, 291

scoparia (*Thelephora*), 222

Sebacina, 190, 197, 198, 199

Senecio, 263; alatipes, 270; albonervius, 275; alienus, 278; alvarezensis, 275; angulifolius, 277, var. **ingens**, 276; anisophyllus, 278; appendiculatus, 265; *arizonicus*, 267; brachyanthus, 277; Breweri, 265; callosus, 271; canus, 266; carnerensis, 273; chapalensis, 277, var. areolatus, 278; Chrismarii, 278; convallium, 266; cordovensis, 277; Coulteri, 272, 273; *diffusus*, 264; *doratophyllus*, 271; Douglasii, 275; *eximius*, 271; *fastigiatus*, 269; filicifolius, 274; Hartwegi, 280; hederæfolius, 278; hyperborealis, 264, var. columbiensis, 264; hypomalacus, 278, *282;* iodanthus, 272, *286;* Kerberi, 279; kernensis, 266; **Klattii**, 281; *latipes*, 271; longilobus, 274; macropus, 267; multilobatus, 265; *neo-mexicanus*, 265, 267, 268; oaxacanus, 279; oreophilus, 267; oreopolus, 268, *284*, forma **aphanactis**, 269; Picridis, 274, 275; præcox, 281; prionopterus, 270; **prolixus**, 264; purpurascens, 273, var. fossanervius, 273; reglensis, 280; *resedifolius*, 264, var. *columbiensis*, 264; roseus, 281; *roseus*, 281; stoechadiformis, 274; subauriculatus, 270, *290;* teliformis, 275; **velatus**, 280, *288; viejensis*, 271; vulgaris, 274; Warszewiczii, 270; Wrightii, 269

Senecioneæ, Descriptions of North American, 263

Septobasidium, 198

Silicic-acid jelly, preparation of, 39

sinuosus (Craterellus), 337

Siphonoglossa, **Greggii**, 409; Pilosella, 410

Sisyrinchium alatum, 407; *alatum* var. *angustissimum*, 406; **angustissimum**, 406, *414*

Skepperia, 343, 345

Solenia, 358, 372; anomala, 373, var. *arbicularis*, 373
spathularius (Craterellus), 345
Sphaeria pocula, 106
spiculosa (Thelephora), 225, *226*
Spirogyra setiformis, isolation of zygospores free from bacteria in, 37
spongiosum (*Stereum*), 215, 216
Staining, A method for the differential, of fungous and host cells, 241
Stemodia, *lanata*, 409; **linearifolia**, 409; **tomentosa**, 409
Stemodiacra linearifolia, 409; *tomentosa*, 409
Stereum, 192, 193, 198, 199; albobadium, destruction of sap-wood of mesquite by, 248; *calyculus*, 338; Leveillianum, destruction of sap-wood of mesquite by, 248; sanguinolentum, latex in, 194; spadiceum, latex in, 194; *spongiosum*, 215, 216
Stichococcus bacillaris, isolation of, in pure culture, 32; relation of, to elementary nitrogen, 178, 179, 181; S. subtilis, isolation of, in pure culture, 32
Stigeoclonium tenue, isolation of, in pure culture, 36
subcyanea (*Cyphella*), 380
subgelatinosa (Cyphella), 370
subgiganteum (Corticium), 197
sulphurea (Cyphella), 360
Syringa vulgaris, trunk disease of, 253

T

taxophilus (Craterellus), 339, *350*
Temperature, effect of: on growth of algæ in pure culture, 180; on acidity of ripening tomato fruits, 239; on cellulose hydrolysis by Sclerotinia cinerea, 311
tephroleuca (*Thelephora*), 213
terrestris (Thelephora), 199, 218, 219, 222, *228*
texensis (Cyphella), 371, *382*
Thelephora, 193, 196, 197, 198, 199; albido-brunnea, 214, 216, *228;* angustata, 205; anthocephala, 202, 203, *226;* biennis, 215, 216; caespitulans, 204, *346;* caryophyllea, 207, 209, *226;* corbiformis, 211; cornucopioides, 205, 212, *346;* cuticularis, 216, *228;* dentosa, 224, *346;* fimbriata, 222, *226;* grisco-zonata, 221, *228; hiscens*, 207; intybacea, 217, 220, *228; laciniata*, 219; lutosa, 216, *346;* **magnispora**, 211, *226;* multipartita, 205, 208, *226; odorifera*, 214; palmata, 201, 203, *226; palmata americana*, 201; pannosa, 207; **perplexa**, 223, *346;* pusiola, 208, var. terrestris, 209; radiata, 210; *Ravenelii*, 207; regularis, 206, 207, *226;* **scissilis**, 204, *226; scoparia*, 222; spiculosa, 225, *226; tephroleuca*, 213; terrestris, 199, 218, 219, 222, *228;* vialis, 213, *228*
Thelephoraceæ of North America, The, I, 185; II, 327; III, 357.
Thompson, C. H., Greenman, J. M. and. Diagnoses of flowering plants, chiefly from the southwestern United States and Mexico, 405
Tiliæ (Cyphella), 364, *382*
Tiliæ (*Peziza*), 364
Tomato fruits, The effect of certain conditions upon the acidity of, 229
Tomato plants, transpiration of, 15
trachychæta (Cyphella), 379
Trametes, 82, 138; *Abietis*, 143; *hispida* 142; *lactea*, 144; malicola, 140, 141; mollis, 141; nivosus, 143; *ohiensis*, 128; Peckii, 140, 142; Pini, 142; rigida, 139, 140, 141; *robiniophila*, 104; *rubescens*, 145; sepium, 139, 140; serialis, 139, 141; suaveolens, 140
Transpiration, The effect of surface films and dusts on the rate of, 1, 351
Tremellodendron, 197, 199
Tulasnella, 198

U

Ulothrix sp., isolation of, in pure culture, 43
Ulva Lactuca, Indications regarding the source of combined nitrogen for, 229
unicolor (Craterellus), 340, 341, 342, *348*

V

vagum (Corticium), 197
Vaucheria sp., isolation of zoospores free from bacteria in, 36
Vaughan, R. E. A method for the differential staining of fungous and host cells, 241
vialis (Thelephora), 213, *228*
villosa (Cyphella), 365, *382*

W

Watering device, automatic, *356;* construction and use of, in transpiration experiments, 351

X

xanthopus (*Merulius*), 336
Xylaria obovata, 225

Z

Zephyranthes **chrysantha**, 406; Eggersiana, 406